

Lab accuracy in a point of care test device.





Substance abuse testing with more substance.



Features & Benefits

- » Integrated test cup screens for a wide variety of prescription and illicit drugs
- » FDA 510(k) Cleared and **CLIA Waived**
- » Self-contained cup is ideal for sending presumptive positive specimens to the lab for confirmation
- » Built-in specimen validity test (SVT) and temperature strip help to monitor specimen tampering (SVT includes BI, Cr, Ni, pH, SG)
- » Fast results in 5 minutes
- » Patented cup design; manufactured in our US facility
- » Long shelf-life

How it works. PRODUCT TRAINING AND CERTIFICATION AVAILABLE ONLINE.

Collect specimen and secure the test lid.




Tilt the cup on its side to activate test.



Interpret test results and specimen validity at 5 minutes.





PRODUCT PROCEDURE NOTE: Refer to product insert for complete instructions, limitations, and warnings.

For in vitro diagnostic use only. This assay provides only a preliminary analytical test result. A more specific alternate chemical method must be used in order to obtain a confirmed analytical result. Gas chromatography/mass spectrometry (GC/MS) is the preferred confirmatory method. Clinical consideration and professional judgement should be applied to any drug of abuse test result, particularly when preliminary positive results are indicated.

Multi-drug configurations

| Item # | iCup [®] Dx with Specimen Validity |
|----------------|---|
| I-DXA-167-013 | BZO, COC, mAMP, OPI, OXY, THC (BI, Cr, Ni, pH, SG) |
| I-DXA-187-016 | AMP, BZO, COC, mAMP, MTD, OPI, OXY, THC (BI, Cr, Ni, pH, SG) |
| I-DXA-1107-142 | AMP, BAR, BZO, COC, mAMP, MDMA, MTD, OPI, OXY, THC (BI, Cr, Ni, pH, SG) |
| I-DXA-1127-023 | AMP, BAR, BZO, COC, mAMP, MDMA, MTD, OPI, OXY, PCP, TCA, THC (BI, Cr, Ni, pH, SG) |

Drugs of abuse abbreviations and cutoffs: Amphetamine (AMP 1000 ng/mL), Barbiturates (BAR 300 ng/mL), Benzodiazepines (BZO 300 ng/mL), Cocaine Metabolites (COC 300 ng/mL), Marijuana (THC 50 ng/mL), MDMA (Ecstasy 500 ng/mL), Methadone (MTD300 ng/mL), Methamphetamine (mAMP 1000 ng/mL), Opiates (OPI 2000 ng/mL), Oxycodone (OXY 100 ng/mL), Phencyclidine (PCP 25 ng/mL), Tricyclic Antidepressants (TCA 1000 ng/mL).

Specimen validity test abbreviations: Bleach (BI), Creatinine (CR), Nitrate (NI), ph (pH), Specific Gravity (SG).

